arXiv:1205.3308v1 [math.NA] 15 May 2012

Convergence Rates and Explicit Error Bounds of Hill's Method for Spectra of Self-Adjoint Differential Operators

Ken'ichiro Tanaka* †, Sunao Murashige* ‡
*School of Systems Information Science, Future University Hakodate,
116-2 Kamedanakano-cho, Hakodate, Hokkaido, 041-8655, Japan
† ketanaka@fun.ac.jp, ‡ murasige@fun.ac.jp

May 15, 2012

Abstract

We present the convergence rates and the explicit error bounds of Hill's method, which is a numerical method for computing the spectra of ordinary differential operators with periodic coefficients. This method approximates the operator by a finite dimensional matrix. On the assumption that the operator is self-adjoint, it is shown that, under some conditions, we can obtain the convergence rates of eigenvalues with respect to the dimension and the explicit error bounds. Numerical examples demonstrate that we can verify these conditions using Gershgorin's theorem for some real problems. Main theorems are proved using the Dunford integrals which project an eigenvector to the corresponding eigenspace.

Keywords Hill's method, Convergence rate, Error bound, Differential Operator, Eigenvalue Problem

Mathematics Subject Classification (2000) 65L15, 65L20, 65L70

1 Introduction

This paper considers Hill's method [8], which is a numerical method for computing spectra of ordinary differential operators S_p with periodic coefficients:

$$S_p = \frac{\mathrm{d}^p}{\mathrm{d}x^p} + \sum_{j=0}^{p-1} \tilde{f}_j(x) \frac{\mathrm{d}^j}{\mathrm{d}x^j}, \tag{1.1}$$

where $x \in \mathbf{R}$, and the functions $\tilde{f}_j$ $(j = 0, 1, \ldots, p-1)$ are C^∞ and satisfy $\tilde{f}_j(x + L) = \tilde{f}_j(x)$ for some real positive constant L. The eigenvalue problem for S_p is described as $S_p\phi = \lambda\phi$, where λ is an eigenvalue and ϕ is an eigenvector. This eigenvalue problem often appears in physical problems such as linear stability analysis of periodic solutions of nonlinear wave equations [7]. The set of eigenvalues λ is included in the spectrum $\sigma(S_p)$. Accordingly, computation of $\sigma(S_p)$ is important in both theoretical and practical points of view. It was reported in [7] that Hill's method can produce very good computed results of $\sigma(S_p)$ for some problems. Note that implementation of Hill's method is straightforward.

The ideas of Hill's method are to employ the Floquet-Bloch decomposition of the spectrum $\sigma(S_p)$ and to approximate the eigenvector ϕ by a finite Fourier series. This Fourier series approximation generates an eigenvalue problem of finite dimension D corresponding to the operator S_p. The aims of this paper are to show the convergence rates of the approximate eigenvalues with respect to the dimension D, and to explicitly obtain the error bounds of them.

On the convergence property of Hill's method, Curtis and Deconinck [4] proved that the exact eigenvalues exist near the computed ones for general cases of S_p, and showed that all exact eigenvalues can be approximated by the computed ones for the case of self-adjoint S_p. Also, they obtained the convergence rate for the case of self-adjoint S_p and the constant coefficients $\tilde{f}_1, \ldots, \tilde{f}_{p-1}$. Johnson and Zumbrun [10] investigated Hill's method using the Evans function, of which the roots correspond to eigenvalues, for general cases of S_p. They showed that the approximate eigenvalues converge to the exact ones, but did not get the convergence rate of them. Vainikko

[11][14] examined an approximation method for the eigenvalue problem $S_p\phi = \lambda\phi$ in an abstract framework, and obtained the convergence rate, which is based on the resolvent norm convergence of approximate operators, for general cases of S_p. We can apply Vainikko's results to Hill's method. But it is difficult to directly evaluate the value of the convergence rate. It should be noted that, although all of these convergence rates yield some error bounds of approximate eigenvalues with unknown coefficients, any explicit error bounds have not been shown.

In this paper, we give a priori estimates of the convergence rate and a posteriori explicit error bounds of Hill's method for self-adjoint S_p with the two cases of coefficient functions $\tilde{f}_j$ in (1.1), namely $\tilde{f}_j \in C^\infty$ and $\tilde{f}_j$ being analytic on some strip region containing the real line. These classes of $\tilde{f}_j$ are more general than [4]. The key ideas of these estimations are to project eigenvectors using the Dunford integrals (5.1) and (5.20), and to specify the disks around the exact eigenvalues using Gershgorin's theorem (Theorem 3.3).

This paper is organized as follows. Section 2 describes Hill's method. In Section 3, we give the convergence rates and the explicit error bounds of Hill's method for self-adjoint differential operators. Section 4 presents numerical examples which supports our results. Section 5 summarizes the proofs of theorems in Section 3. Section 6 concludes this paper.

2 Hill's method

This section describes Hill's method [8] which is a numerical method for computing the spectum $\sigma(S_p)$ of the ordinary differential operator S_p with periodic coefficients $\tilde{f}_j$ defined by (1.1). This operator S_p can be regarded as an operator $S_p : H^p(\mathbf{R}) \to L_2(\mathbf{R})$, where $L_2(\mathbf{R})$ is the Lebesgue space of square integrable functions on $\mathbf{R}$ and $H^p(\mathbf{R}) \subset L_2(\mathbf{R})$ is the Sobolev space of functions whose derivatives up to p-th order are square integrable. Hill's method approximates the elements of $\sigma(S_p)$ by the following two steps.

Step 1: Floquet-Bloch decomposition. In order to apply the Floquet theory, we introduce a new operator $S_p^\mu : H^p([0, L])_{\mathrm{per}} \to L_2([0, L])_{\mathrm{per}}$ defined by

$$S_p^\mu = \mathrm{e}^{-\mathrm{i}\mu x} S_p \mathrm{e}^{\mathrm{i}\mu x}, \tag{2.1}$$

where $\mu \in [0,\, 2\pi/L)$, and $L_2([0, L])_{\mathrm{per}}$ and $H^p([0, L])_{\mathrm{per}}$ are the Lebesgue space and the Sobolev space of periodic functions on $[0, L]$, respectively. Note that S_p is defined on $\mathbf{R}$, whereas S_p^μ is on $[0, L]$. If S_p is self-adjoint, then S_p^μ is also self-adjoint. Moreover, S_p^μ is explicitly written as

$$S_p^\mu = \frac{\mathrm{d}^p}{\mathrm{d}x^p} + \sum_{j=0}^{p-1} f_j(x) \frac{\mathrm{d}^j}{\mathrm{d}x^j} \tag{2.2}$$

for some periodic functions $f_0, \ldots, f_{p-1}$ with period L.

It is known that the spectrum $\sigma(S_p^\mu)$ consists of only the eigenvalues of S_p^μ, and the Floquet theory yields

$$\sigma(S_p) = \bigcup_{\mu \in [0,\, 2\pi/L)} \sigma(S_p^\mu). \tag{2.3}$$

This decomposition (2.3) is called the Floquet-Bloch decomposition. Accordingly, it suffices to consider the eigenvalue problem $S_p^\mu \phi = \lambda\phi$ for $\mu \in [0,\, 2\pi/L)$, where $\lambda \in \mathbf{C}$ is an eigenvalue and $\phi \in L_2([0, L])_{\mathrm{per}}$ is an eigenvector. In addition, it should be noted that $\sigma(S_p^\mu)$ is a discrete set of the eigenvalues of S_p^μ without accumulation points. This discreteness follows from the compactness of the resolvent of S_p^μ provided $\rho(S_p^\mu) \neq \emptyset$, where $\rho(S_p^\mu) = \mathbf{C} \setminus \sigma(S_p^\mu)$. See e.g. Lemma 2 in [6, Chapter XIX Section 2], Lemma 3 in [6, Chapter XIX Section 3] and Lemma 16 in [5, Chapter XIII Section 2].

Step 2: Fourier series approximation. Since $\phi \in L_2([0, L])_{\mathrm{per}}$ has the Fourier series expansion:

$$\phi(x) = \frac{1}{\sqrt{L}} \sum_{n=-\infty}^{\infty} \hat{\phi}_n \exp\left(-\mathrm{i}\frac{2\pi n x}{L}\right), \tag{2.4}$$

ϕ can be approximated by the truncation of this series:

$$(\hat{P}_N \phi)(x) = \frac{1}{\sqrt{L}} \sum_{n=-N}^{N} \hat{\phi}_n \exp\left(-\mathrm{i}\frac{2\pi n x}{L}\right). \tag{2.5}$$

This truncation reduces the eigenvalue problem $S_p^\mu \phi = \lambda \phi$ to a finite dimensional problem. More precisely, the problem

$$S_{p,N}^\mu \phi_N = \lambda_N \phi_N \quad \text{with} \quad S_{p,N}^\mu = \hat{P}_N S_p^\mu \hat{P}_N \tag{2.6}$$

gives approximate eigenvalues λ_N's for the original problem $S_p^\mu \phi = \lambda \phi$.

Since the problem (2.6) is equivalent to a matrix eigenvalue problem, we can obtain the approximate values of the eigenvalues of S_p^μ using some standard numerical method. In the following sections, let $\sigma(S_{p,N}^\mu)$ denote the set of eigenvalues of $S_{p,N}^\mu$.

3 Convergence Rates and Error Bounds of Hill's Method

In this section, we present theorems about a priori estimates of the convergence rate and a posteriori explicit error bounds of Hill's method for the eigenvalue problem $S_p^\mu \phi = \lambda \phi$ with self-adjoint S_p^μ on some assumptions. Their proofs are given in Section 5. In what follows, we use the notations defined in Section 2.

3.1 Assumptions

First, we assume the self-adjointness of the operator S_p^μ.

Assumption 1. *The operator S_p^μ is self-adjoint, i.e. $(S_p^\mu)^* = S_p^\mu$, where * represents the adjoint of operators.*

On Assumption 1, $S_{p,N}^\mu$ in (2.6) is also self-adjoint.

Next, we specify the smoothness of the coefficient functions f_j's of S_p^μ in (2.2).

Assumption 2a. *For the operator S_p^μ, the coefficients $f_0, f_1, \ldots, f_{p-1}$ are C^∞.*

In addition, as a special case of Assumption 2a, the following assumption is prepared.

Assumption 2b. *For the operator S_p^μ, the coefficients $f_0, f_1, \ldots, f_{p-1}$ are analytic on a complex domain*

$$\mathcal{D}_d = \{z \in \mathbf{C} \mid |\mathrm{Im}\, z| < d\} \tag{3.1}$$

for some $d > 0$.

Assumption 2b is often satisfied in real problems.

If $S_{p,N}^\mu$ approximates S_p^μ very well, we may expect that approximate eigenvalues λ_N's for large enough N should be included in the neighborhood $B_\lambda(r_\lambda)$ of the corresponding exact eigenvalue λ, where

$$B_\zeta(r) = \{z \in \mathbf{C} \mid |z - \zeta| \le r\}. \tag{3.2}$$

Then we prepare the following two assumptions.

Assumption 3a. *For $\lambda \in \sigma(S_p^\mu)$, there exists a positive real number r_λ such that the following holds true: for some sequence $\{\lambda_N \mid \lambda_N \in \sigma(S_{p,N}^\mu)\}$ and some positive integer N_0, if $N > N_0$ then*

$$|\lambda - \lambda_N| \le \frac{r_\lambda}{2} \tag{3.3}$$

and

$$2r_\lambda \le \min\left\{\mathrm{dist}(\lambda, \sigma(S_p^\mu) \setminus \{\lambda\}),\, \mathrm{dist}(\lambda, \sigma(S_{p,N}^\mu) \setminus \{\lambda_N\})\right\}. \tag{3.4}$$


$\Gamma_\lambda(S_p^\mu)$
$r_\lambda/2$
r_λ
$2r_\lambda$
λ
λ_N
Elements of
$\sigma(S_p^\mu) \setminus \{\lambda\}$ or
$\sigma(S_{p,N}^\mu) \setminus \{\lambda_N\}$

Figure 1: Illustration of the conditions (3.3) and (3.4) in Assumption 3a. This figure depicts locations of an exact eigenvalue $\lambda \in \sigma(S_p^\mu)$, the corresponding approximate eigenvalue $\lambda_N \in \sigma(S_{p,N}^\mu)$ and some other eigenvalues on the complex plane $\mathbf{C}$. The eigenvalue λ_N is located inside the circle with center λ and radius $r_\lambda/2$ (shown by the smaller dotted circle), and the other eigenvalues than λ and λ_N are outside the circle with center λ and radius $2r_\lambda$ (shown by the larger dotted circle). The directed contour $\Gamma_\lambda(S_p^\mu)$ is used to prove the theorems in Sections 3.2 and 3.3.

Figure 1 illustrates these conditions (3.3) and (3.4). When a single sequence $\{\lambda_N\}$ for some $\lambda \in \sigma(S_p^\mu)$ is detected, Assumption 3a is effective for estimation of the convergence rate and the error bound of $\{\lambda_N\}$. In general, there exist more than one sequence of approximate eigenvalues for some $\lambda \in \sigma(S_p^\mu)$. Then we set another assumption as follows:

Assumption 3b. *Let k be an integer with $k \geq 2$. For $\lambda \in \sigma(S_p^\mu)$, there exists a positive real number r_λ such that the following holds true: for some sequences $\{\lambda_{N,i} \mid \lambda_{N,i} \in \sigma(S_{p,N}^\mu)\}$ $(i = 1, 2, \dots k)$ and some positive integer N_0, if $N > N_0$ then*

$$|\lambda - \lambda_{N,i}| \leq \frac{r_\lambda}{2} \quad (i = 1, 2, \dots, k) \tag{3.5}$$

and

$$2r_\lambda \leq \min \left\{ \mathrm{dist}(\lambda, \sigma(S_p^\mu) \setminus \{\lambda\}),\ \mathrm{dist}\left(\lambda,\ \sigma(S_{p,N}^\mu) \setminus \left(\bigcup_{i=1}^{k} \{\lambda_{N,i}\} \right)\right) \right\}. \tag{3.6}$$

Remark 3.1. *Assumptions 3a and 3b mean that, for sufficiently large N, the number of eigenvalues of $S_{p,N}^\mu$ approximating $\lambda \in \sigma(S_p^\mu)$ is at most finite, and the other elements of $\sigma(S_p^\mu)$ and $\sigma(S_{p,N}^\mu)$ are relatively far from λ. The finiteness of the approximate eigenvalues is based on the fact that the eigenspace of S_p^μ for any λ is finite dimensional.*

When Assumption 3a or 3b are satisfied, there exist some convergent subsequences in $\bigcup_N \sigma(S_{p,N}^\mu)$. Thus these assumptions are closely related to the convergence property of the eigenvalues of $S_{p,N}^\mu$. On this convergence property, the following two theorems are known [4][12]. Theorem 3.1 gives so-called "no-spurious-mode condition".

Theorem 3.1 ([4, Theorem 9])**.** *Let $\mathfrak{D} \subset \mathbf{C}$ be a compact set and $\{\lambda_N\}$ be a sequence in $\mathfrak{D}$ with $\lambda_N \in \sigma(S_{p,N}^\mu)$. Then for any $\varepsilon > 0$ there exists a positive integer M such that the following holds true: for any N with $N \geq M$, λ_N is contained in the ε-neighborhood of some $\lambda \in \mathfrak{D} \cap \sigma(S_p^\mu)$.*

Theorem 3.2 ([4, Corollary of Lemma 10 and Theorem 16])**.** *Assume that S_p^μ is self-adjoint. Then for any $\lambda \in \sigma(S_p^\mu)$ there exists some subsequence $\{\lambda_{N_j}\}$ of $\{\lambda_N\}$ with $\lambda_N \in \sigma(S_{p,N}^\mu)$ such that $\lambda_{N_j} \to \lambda$.*

In Section 3.3, a sufficient condition for (3.3) and (3.4) in Assumption 3a is given by Theorem 3.6. We can verify this sufficient condition using Theorems 3.1, 3.2 and the following theorem for a concrete problem as shown in Section 4.

Theorem 3.3 (Gershgorin's Theorem [15]). *For any $n \times n$ matrix $V = (v_{ij})$ with $v_{ij} \in \mathbf{C}$, all eigenvalues of V are contained in $\bigcup_{i=1}^{n} C_i$, where $C_1, \ldots, C_n$ are disks defined as*

$$C_i = \left\{ z \in \mathbf{C} \;\middle|\; |z - v_{ii}| \leq \sum_{j \neq i} |v_{ij}| \right\}. \tag{3.7}$$

Moreover, each connected component of $\bigcup_{i=1}^{n} C_i$ contains as many eigenvalues of V as the disks composing it.

This theorem is used to estimate the radius r_λ in the conditions (3.3) and (3.4), whereas Theorems 3.1 and 3.2 guarantee that the sequence of the approximate eigenvalues converges to an exact eigenvalue.

3.2 A Priori Estimates of Convergence Rates

When Assumptions 1, 2 and 3 are satisfied, we can obtain a priori estimates of the convergence rates as follows.

Theorem 3.4. *Let Assumptions 1 and 3a be satisfied. Then, when Assumption 2a is satisfied, namely f_j being C^∞, for a sufficiently large N and any positive integer q, there exists a positive constant C_1 such that*

$$|\lambda - \lambda_N| \leq C_1\, N^{-q}, \tag{3.8}$$

where C_1 depends only on p, $f_0, \ldots, f_{p-1}$, λ, r_λ, q and an eigenvector ϕ of S_p^μ corresponding to λ. In addition, when Assumption 2b is satisfied, namely f_j being analytic, for a sufficiently large N and any ε with $0 < \varepsilon < d$, there exists a positive constant C_2 such that

$$|\lambda - \lambda_N| \leq C_2\, N^{p+1/2} \exp\left(-\frac{2\pi(d-\varepsilon)}{L} N\right), \tag{3.9}$$

where C_2 depends only on p, $f_0, \ldots, f_{p-1}$, λ, r_λ, d, ε and ϕ.

Theorem 3.5. *Let Assumptions 1 and 3b be satisfied. Then, when Assumption 2a is satisfied, namely f_j being C^∞, for a sufficiently large N, some choice of an index $i_N \in \{1, \ldots, k\}$ and any positive integer q, there exists a positive constant C_1 such that*

$$|\lambda - \lambda_{N,i_N}| \leq C_1\, N^{-q}, \tag{3.10}$$

where C_1 depends only on p, $f_0, \ldots, f_{p-1}$, λ, r_λ, q and an eigenvector ϕ of S_p^μ corresponding to λ. In addition, when Assumption 2b is satisfied, namely f_j being analytic, for a sufficiently large N, some choice of an index $i_N \in \{1, \ldots, k\}$ and any ε with $0 < \varepsilon < d$, there exists a positive constant C_2 such that

$$|\lambda - \lambda_{N,i_N}| \leq C_2\, N^{2p+1} \exp\left(-\frac{2\pi(d-\varepsilon)}{kL} N\right), \tag{3.11}$$

where C_2 depends only on p, $f_0, \ldots, f_{p-1}$, λ, r_λ, d, ε and ϕ.

3.3 A Posteriori Error Bounds

When Assumptions 1 and 2a are satisfied, we can explicitly obtain a posteriori estimates of the error bounds as follows. Here, for simplicity, only the case of Assumption 3a is considered.

Theorem 3.6. *Let Assumptions 1 and 2a be satisfied. Assume that there exists $\zeta \in \mathbf{C}$ and $r > 0$ such that, for any N,*

$$\lambda_N \in B_\zeta(r) \quad \text{and} \quad B_\zeta(9r) \cap \left(\sigma(S_{p,N}^\mu) \setminus \{\lambda_N\}\right) = \emptyset, \tag{3.12}$$

where $\lambda_N \in \sigma(S_{p,N}^\mu)$ and $B_\zeta(r)$ is defined by (3.2). Then there uniquely exists $\lambda \in \sigma(S_p^\mu) \cap B_\zeta(r)$ satisfying

$$\begin{aligned} |\lambda - \lambda_N| \leq{}& \left(5 + \frac{3|\zeta|}{r}\right) \frac{(2\pi N)^p}{L^{p+1/2}} \\ &\cdot \sum_{j=0}^{p} \left(\sum_{N<|l|<2N} \sum_{m=l-N}^{l+N} \left| (\hat{f}_j)_m (\hat{\phi}_N)_{l-m} \right| + (2N+1) \sum_{|m| \geq N} \left| (\hat{f}_j)_m \right| \right), \end{aligned} \tag{3.13}$$

where $f_p \equiv 1$, ϕ_N *is an eigenvector corresponding to* λ_N, *and* $(\hat{f}_j)_m$ $(m = 0, \pm 1, \pm 2, \ldots)$ *and* $(\hat{\phi}_N)_n$ $(n = 0, \pm 1, \pm 2, \ldots)$ *are the Fourier coefficients of* f_j $(j = 0, \ldots, p)$ *and* ϕ_N, *respectively.*

4 Numerical Experiments

In this section, we apply Hill's method described in Section 2 to Hill's operator (4.1) of which the spectrum is exactly known, and observe that computed results are consistent with Theorems 3.4, 3.5 and 3.6 in Section 3.

4.1 Hill's Operator

As an example for numerical experiments, we consider Hill's operator [7] defined by

$$S_2 = -\frac{\mathrm{d}^2}{\mathrm{d}x^2} + \{6\ell^2 \operatorname{sn}^2(x, \ell) - 4 - \ell^2\}, \tag{4.1}$$

where $\operatorname{sn}(\cdot, \ell)$ is the Jacobian elliptic function with modulus ℓ $(0 \leq \ell < 1)$. It is known [7] that the spectrum $\sigma(S_2)$ is exactly given by

$$\sigma(S_2) = [\sigma_{\mathrm{a}}(\ell),\, -3] \cup [\sigma_{\mathrm{b}}(\ell),\, 0] \cup [\sigma_{\mathrm{c}}(\ell),\, +\infty), \tag{4.2}$$

where

$$\begin{aligned} &\text{(a)} \quad \sigma_{\mathrm{a}}(\ell) = \ell^2 - 2 - 2\sqrt{1 - \ell^2 + \ell^4}, \\ &\text{(b)} \quad \sigma_{\mathrm{b}}(\ell) = -3(1 - \ell^2), \\ &\text{(c)} \quad \sigma_{\mathrm{c}}(\ell) = \ell^2 - 2 + 2\sqrt{1 - \ell^2 + \ell^4}. \end{aligned} \tag{4.3}$$

Figure 2 illustrates $\sigma(S_2)$ for all ℓ with $0 \leq \ell < 1$.

By definition (2.2) of S_p^μ, $p = 2$ and S_2^μ is expressed as

$$S_2^\mu = -\frac{\mathrm{d}^2}{\mathrm{d}x^2} + f_1(x)\frac{\mathrm{d}}{\mathrm{d}x} + f_0(x) \tag{4.4}$$

with

$$f_1(x) = -2\mathrm{i}\mu \quad \text{and} \quad f_0(x) = \{6\ell^2 \operatorname{sn}^2(x, \ell) - 4 - \ell^2\} + \mu^2. \tag{4.5}$$

Although the sign of the highest order derivative $\mathrm{d}^2/\mathrm{d}x^2$ in (4.4) is different from that in (2.1), we can directly apply the results in Section 3 to S_2^μ. Note that f_0 is periodic with period $2K(\ell)$, where $K(\ell)$ is the complete elliptic integral of the first kind with modulus ℓ. Since S_2^μ is self-adjoint and f_0 is analytic with $d = K\left(\sqrt{1 - \ell^2}\right)$ in (3.1), Assumptions 1 and 2b are satisfied.

For later convenience, expand the periodic term $f_0(x) - \mu^2$ in the Fourier series form

$$f_0(x) - \mu^2 = b_0 + \sum_{j=1}^{\infty} b_j \left\{ \exp\left(\mathrm{i}\frac{\pi j x}{K(\ell)}\right) + \exp\left(-\mathrm{i}\frac{\pi j x}{K(\ell)}\right) \right\},$$

where

$$b_0(\ell) = 6\left(1 - \frac{E(\ell)}{K(\ell)}\right) - 4 - \ell^2, \quad b_j(\ell) = -\frac{6\pi^2}{K(\ell)^2} \frac{j\, q(\ell)^j}{1 - q(\ell)^{2j}} \quad (j = 1, 2, \ldots), \tag{4.6}$$

$E(\ell)$ is the complete elliptic integral of the second kind with modulus ℓ, and

$$q(\ell) = \exp\left(-\pi \frac{K\left(\sqrt{1 - \ell^2}\right)}{K(\ell)}\right). \tag{4.7}$$



Figure 2: Computed results of $(\sigma(S_2), \ell)$ for S_2 in (4.1) (a) $\sigma_{\mathrm{a}}(\ell) = \ell^2 - 2 - 2\sqrt{1 - \ell^2 + \ell^4}$, (b) $\sigma_{\mathrm{b}}(\ell) = -3(1 - \ell^2)$ and (c) $\sigma_{\mathrm{c}}(\ell) = \ell^2 - 2 + 2\sqrt{1 - \ell^2 + \ell^4}$.

Then, for $L = 2MK(\ell)$ with positive integer M, the eigenvalue problem $S^{\mu}_{2,N}\phi = \lambda\phi$ for $\phi \in L_2([0, L])_{\mathrm{per}}$ can be written in the form

$$\hat{D}^{\mu}_{n}(\ell)\ (\hat{\phi}_N)_n + \sum_{j=1}^{N} b_j(\ell) \Big((\hat{\phi}_N)_{n-Mj} + (\hat{\phi}_N)_{n+Mj} \Big) = \lambda(\hat{\phi}_N)_n \quad (-N \le n \le N) \tag{4.8}$$

with

$$\hat{D}^{\mu}_{n}(\ell) = \left(\mu + \frac{\pi n}{MK(\ell)}\right)^2 + b_0(\ell) \quad (n = -N, \ldots, N). \tag{4.9}$$

Equation (4.8) is a matrix eigenvalue problem of $2N + 1$ dimension.

For given M and N, we can compute the eigenvalues of $S^{\mu}_{2,N}$ using Hill's method. Figure 3 shows the absolute errors of the computed eigenvalues corresponding to (a) in (4.3) for $M = 2$ and $N = 5, 10, \ldots, 40$. Here it should be noted that the boundaries (a), (b) and (c) in (4.3) are obtained as the eigenvalues of S^{μ}_{p} in (4.4) for $\mu = 0$. All programs used in this paper are written in C with quadruple-precision floating-point arithmetic and executed on a computer with the Sparc processor and the Fujitsu compiler C99.

4.2 The Convergence Rate

The computed eigenvalues of $S^{0}_{2,N}$ approach to the exact ones corresponding to them with increase of N as shown in Figure 3. Thus the conditions (3.3) and (3.4) in Assumption 3a are satisfied for sufficiently large N, and the convergence rate (3.9) in Theorem 3.4 can be applied to the computed eigenvalues λ_N's. In fact, the exponential decay of the error with N in (3.9) can be found in Figure 3. Note that similar results to Figure 3 are obtained for the eigenvalues (b) and (c). From these, we can say that the computed results are consistent with (3.9) in Theorem 3.4.

For small values of ℓ, we can show that the convergence rate (3.9) holds, even if the exact eigenvalues of S^{0}_{2} are unknown, as follows. For that, first, consider (3.12) which is the sufficient condition for (3.3) and (3.4) in Assumption 3a. We can check (3.12) by applying Theorem 3.3 (Gershgorin's theorem) to the coefficient matrix in the left hand side of (4.8), of which the diagonal elements are given by $\hat{D}^{\mu}_{n}(\ell)$ in (4.9) and the eigenvalues are real. For this matrix, Theorem 3.3 produces

$$\sigma(S^{\mu}_{2,N}) \subset \bigcup_{n=-N}^{N} I^{\mu}_{n}(\ell) \tag{4.10}$$

where

$$I_n^\mu(\ell) = \left\{ z \in \mathbf{R} \,\middle|\, |z - \hat{D}_n^\mu(\ell)| \le r(\ell) \right\} \tag{4.11}$$

with

$$r(\ell) = \frac{12\pi^2}{K(\ell)^2} \frac{q(\ell)}{(1-q(\ell)^2)(1-q(\ell))^2}. \tag{4.12}$$

We can compute the center $\hat{D}_n^\mu(\ell)$ in (4.9) and the radius $r(\ell)$ in (4.12) of the interval $I_n^\mu(\ell)$ which includes some eigenvalues in $\sigma(S_{2,N}^\mu)$. For example, when $\ell = 0.1$ and $\mu = 0$, we have

$$r(0.1) = 0.030\cdots,$$

and

$$\begin{aligned} &\hat{D}_0^0(0.1) = -3.9799\cdots, \quad \hat{D}_{\pm 1}^0(0.1) = -2.9849\cdots, \\ &\hat{D}_{\pm 2}^0(0.1) \doteqdot 0, \quad \hat{D}_{\pm 3}^0(0.1) = 4.9749\cdots. \end{aligned}$$

Then the intervals $I_n^0(0.1)$ $(n = 0, 1, 2, 3)$ are mutually disjoint and the distances between their centers are larger than $10\, r(0.1)$. Moreover, from Theorem 3.3, we can say that $I_0^0(0.1)$ contains just one eigenvalue of $S_{2,N}^\mu$. Note that ζ and r in (3.12) correspond to $\hat{D}_0^0(0.1)$ and $r(0.1)$, respectively. Therefore the condition (3.12) is satisfied for the interval $I_0^0(0.1)$. From these, the conditions (3.3) and (3.4) in Assumption 3a are satisfied, and the convergence rate (3.9) holds for $I_0^0(0.1)$.

Also from Theorem 3.3, we can show that $I_1^0(0.1)$ and $I_2^0(0.1)$ contain just two eigenvalues of $S_{2,N}^\mu$, respectively. Then the conditions (3.5) and (3.6) in Assumption 3b may be satisfied, and it was found that the numerical results are consistent with the convergence rate (3.11). For further study on this case of Assumption 3b, we have to develop the verification method for the conditions (3.5) and (3.6).

4.3 The Error Bound

Since the computed eigenvalues are accurate enough for sufficiently large N, as shown in Figure 3, the condition (3.12) in Theorem 3.6 is satisfied for sufficiently large N. Then we can estimate the error bound (3.13) in Theorem 3.6 as follows. First, ζ, r, L and $(\hat{f}_0)_m$ in (3.13) are given by, respectively,

$$\zeta = \hat{D}_0^0(\ell),\ r = r(\ell),\ L = 4K(\ell),\ \text{and}\ (\hat{f}_0)_m = \begin{cases} 0 & (m \text{ is odd}), \\ \sqrt{L}\, b_{m/2} & (m \text{ is even}), \end{cases} \tag{4.13}$$

for $|m| \ge 1$. Next, the infinite sum in (3.13) can be bounded as

$$\begin{aligned} \sum_{|m| \ge N} \left| (\hat{f}_0)_m \right| &\le 2\sqrt{L} \sum_{j=\lceil N/2 \rceil}^{\infty} |b_j| \le \frac{24\pi^2}{K(\ell)^{3/2}} \sum_{j=\lceil N/2 \rceil}^{\infty} \frac{j\, q(\ell)^j}{1-q(\ell)^2} \\ &= \frac{24\pi^2}{K(\ell)^{3/2}} \frac{(\lceil N/2 \rceil (1-q(\ell)) + q(\ell))\, q(\ell)^{\lceil N/2 \rceil}}{(1-q(\ell)^2)(1-q(\ell))^2}. \end{aligned} \tag{4.14}$$

Then it follows from the inequality (3.13) that

$$\begin{aligned} |\lambda - \lambda_N| \le \left(5 + \frac{3|\zeta|}{r}\right) \frac{(2\pi N)^p}{L^{p+1/2}} \Bigg(&\sum_{N<|l|<2N} \sum_{m=l-N}^{l+N} \left| (\hat{f}_0)_m (\hat{\phi}_N)_{l-m} \right| \\ &+ (2N+1) \frac{24\pi^2}{K(\ell)^{3/2}} \frac{(\lceil N/2 \rceil (1-q(\ell)) + q(\ell))\, q(\ell)^{\lceil N/2 \rceil}}{(1-q(\ell)^2)(1-q(\ell))^2} \Bigg), \end{aligned} \tag{4.15}$$

where ζ, r, L and $(\hat{f}_0)_m$ are given by (4.13). The computed results of the right hand side of (4.15) are shown in Figure 4. The absolute errors in Figure 3 are bounded by the corresponding error bounds in Figure 4. Thus the numerical results are consistent with Theorem 3.6.



Figure 3: The absolute errors of the approximate eigenvalues of $\sigma(S^0_{2,N})$ corresponding to the eigenvalues (a).



Figure 4: The error bounds (4.15) of the approximate eigenvalues of $\sigma(S^0_{2,N})$ corresponding to the eigenvalues (a).

For small values of ℓ, we can obtain the explicit error bound (4.15), even if the exact eigenvalues of S^0_2 are unknown. For example, when $\ell = 0.1$ and $\mu = 0$, the interval $I^0_0(0.1)$ in (4.11) satisfies the condition (3.12) as shown in Section 4.2. Then we can conclude that $I^0_0(0.1)$ contains one exact eigenvalue of S^0_2, and the corresponding error is bounded by (4.15). Figure 4 shows that, when $\ell = 0.1$, the error for $N \geq 10$ is less than 2.37×10^{-8}.

5 Proofs

This section summarizes proofs of the theorems in Sections 3.2 and 3.3. Let $(\cdot\,,\,\cdot)_2$ and $\|\cdot\|_2$ be the inner product of $L_2([0,L])_{\mathrm{per}}$ and the norm induced by it, respectively. We also use $\|\cdot\|_2$ as the operator norm of linear operators on $L_2([0,L])_{\mathrm{per}}$ with respect to the norm $\|\cdot\|_2$.

5.1 Proof of Theorem 3.4

To prove the theorem, we use an appropriate eigenvector ϕ_N of the approximate operator $S^\mu_{p,N}$ generated by an eigenvector ϕ of S^μ_p. On Assumption 3a, we can employ the projection operator $P_\lambda(S^\mu_{p,N})$ to the eigenspace corresponding to $\lambda_N \in \sigma(S^\mu_{p,N})$ defined by

$$P_\lambda(S^\mu_{p,N}) = \frac{1}{2\pi \mathrm{i}} \oint_{\Gamma_\lambda(S^\mu_p)} R_\nu(S^\mu_{p,N})\,\mathrm{d}\nu, \tag{5.1}$$

where $R_\nu(S^\mu_{p,N}) = (\nu I - S^\mu_{p,N})^{-1}$ and $\Gamma_\lambda(S^\mu_p)$ is the directed circle with center λ and radius r_λ shown by Figure 1. The integral in (5.1) is the Dunford integral. See e.g. Theorem XII.5 in [13]. For later use, it should be noted that

$$l(\Gamma_\lambda(S^\mu_p)) = 2\pi r_\lambda, \tag{5.2}$$

where $l(\Gamma_\lambda(S^\mu_p))$ is the length of $\Gamma_\lambda(S^\mu_p)$, and

$$|\nu - \lambda| = r_\lambda, \tag{5.3}$$

$$r_\lambda/2 \leq |\nu - \lambda_N| \leq 3r_\lambda/2. \tag{5.4}$$

for any $\nu \in \Gamma_\lambda(S^\mu_p)$.

Then we take an eigenvector ϕ with $\|\phi\|_2 = 1$ corresponding to λ, and using ϕ and the projection $P_{\lambda_N}(S^\mu_{p,N})$ for $\lambda_N \in \sigma(S^\mu_{p,N})$, we can obtain an eigenvector ϕ_N with $\|\phi_N\|_2 = 1$ corresponding to λ_N. The feasibility of such procedure is guaranteed by the following lemma, which is based on the idea in [1].

Lemma 5.1. *Let S_p^μ satisfy Assumptions 1 and 2a, $\lambda \in \sigma(S_p^\mu)$ and $\lambda_N \in \sigma(S_{p,N}^\mu)$ satisfy Assumption 3a. Then for an eigenvector ϕ corresponding to λ we have*

$$\|\phi - P_{\lambda_N}(S_{p,N}^\mu)\phi\|_2 \le \frac{2}{r_\lambda}\|(S_p^\mu - S_{p,N}^\mu)\phi\|_2. \tag{5.5}$$

Furthermore, if $\|\phi\|_2 = 1$ and there exists a positive integer N_0 such that for any $N > N_0$

$$\frac{2}{r_\lambda}\|(S_p^\mu - S_{p,N}^\mu)\phi\|_2 < 1, \tag{5.6}$$

we have $P_{\lambda_N}(S_{p,N}^\mu)\phi \neq 0$ for $N > N_0$.

Proof. Under Assumption 2a, we can use the contour $\varGamma_\lambda(S_p^\mu)$ to define $P_\lambda(S_p^\mu)$ and $P_{\lambda_N}(S_{p,N}^\mu)$. By the residue theorem, we have $P_\lambda(S_p^\mu)\phi = \phi$. Then it follows from (5.2) that

$$\begin{aligned}\|\phi - P_{\lambda_N}(S_{p,N}^\mu)\phi\|_2 &= \|P_\lambda(S_p^\mu)\phi - P_{\lambda_N}(S_{p,N}^\mu)\phi\|_2 \\ &= \left\|\frac{1}{2\pi\mathrm{i}}\oint_{\varGamma_\lambda(S_p^\mu)}(R_\nu(S_p^\mu) - R_\nu(S_{p,N}^\mu))\phi\,\mathrm{d}\nu\right\|_2 \\ &\le r_\lambda \max_{\nu\in\varGamma_\lambda(S_p^\mu)}\|(R_\nu(S_p^\mu) - R_\nu(S_{p,N}^\mu))\phi\|_2. \end{aligned}\tag{5.7}$$

As for $\|(R_\nu(S_p^\mu) - R_\nu(S_{p,N}^\mu))\phi\|_2$ in (5.7), we have

$$\begin{aligned}\|(R_\nu(S_p^\mu) - R_\nu(S_{p,N}^\mu))\phi\|_2 &= \|(\nu I - S_{p,N}^\mu)^{-1}(S_p^\mu - S_{p,N}^\mu)(\nu I - S_p^\mu)^{-1}\phi\|_2 \\ &= \|(\nu I - S_{p,N}^\mu)^{-1}(S_p^\mu - S_{p,N}^\mu)(\nu - \lambda)^{-1}\phi\|_2 \\ &\le |\nu-\lambda|^{-1}\,\|(\nu I - S_{p,N}^\mu)^{-1}\|_2\,\|(S_p^\mu - S_{p,N}^\mu)\phi\|_2 \\ &= |\nu-\lambda|^{-1}\,\mathrm{dist}(\nu,\sigma(S_{p,N}^\mu))^{-1}\,\|(S_p^\mu - S_{p,N}^\mu)\phi\|_2, \end{aligned}\tag{5.8}$$

where $\mathrm{dist}(\nu,\sigma(S_{p,N}^\mu))$ is the distance between $\nu \in \mathbf{C}$ and $\sigma(S_{p,N}^\mu)$ defined as

$$\mathrm{dist}(\nu,\sigma(S_{p,N}^\mu)) = \inf_{\xi\in\sigma(S_{p,N}^\mu)}|\nu-\xi|.$$

The last equality in (5.8) follows from the self-adjointness of $S_{p,N}^\mu$. Furthermore, by (5.3) and (5.4) we have

$$|\nu - \lambda| = r_\lambda, \tag{5.9}$$

for any $\nu \in \varGamma_\lambda(S_p^\mu)$ and

$$\min_{\nu\in\varGamma_\lambda(S_p^\mu)}\mathrm{dist}(\nu,\sigma(S_{p,N}^\mu)) = \min_{\nu\in\varGamma_\lambda(S_p^\mu)}|\nu-\lambda_N| \ge r_\lambda/2. \tag{5.10}$$

Combining (5.8), (5.9) and (5.10) we have

$$\max_{\nu\in\varGamma_\lambda(S_p^\mu)}\|(R_\nu(S_p^\mu) - R_\nu(S_{p,N}^\mu))\phi\|_2 \le \frac{2}{r_\lambda^2}\|(S_p^\mu - S_{p,N}^\mu)\phi\|_2. \tag{5.11}$$

Then we can deduce the conclusion (5.5) from (5.7) and (5.11).

Finally, if $\|\phi\|_2 = 1$ and (5.6) holds for $N > N_0$, it follows from (5.5) that $\|\phi - P_{\lambda_N}(S_{p,N}^\mu)\phi\|_2 < 1$. Hence $P_{\lambda_N}(S_{p,N}^\mu)\phi \neq 0$ holds true. □

Here we present a key lemma for the estimate of the convergence rate. Part of this lemma is also based on the idea in [1].

Lemma 5.2. *Let S_p^μ satisfy Assumptions 1 and 2a, $\lambda \in \sigma(S_p^\mu)$ and $\lambda_N \in \sigma(S_{p,N}^\mu)$ satisfy Assumption 3a. Furthermore, for an eigenvector ϕ corresponding to λ with $\|\phi\|_2 = 1$, assume that there exists a positive integer N_0 such that for any $N > N_0$*

$$\frac{2}{r_\lambda}\|(S_p^\mu - S_{p,N}^\mu)\phi\|_2 < 1. \tag{5.12}$$

Then for $N > N_0$ we have

$$|\lambda - \lambda_N| \leq \left(3 + \frac{8|\lambda|}{r_\lambda}\right)\|(S_p^\mu - S_{p,N}^\mu)\phi\|_2. \tag{5.13}$$

Proof. By Lemma 5.1, for $N > N_0$ we can define

$$\phi_N = \frac{1}{\|P_{\lambda_N}(S_{p,N}^\mu)\phi\|_2} P_{\lambda_N}(S_{p,N}^\mu)\phi. \tag{5.14}$$

Then ϕ_N is an eigenvector corresponding to λ_N with $\|\phi_N\|_2 = 1$. Then noting $\lambda = (S_p^\mu\phi, \phi)_2$ and $\lambda_N = (S_{p,N}^\mu\phi_N, \phi_N)_2$ we have

$$\begin{aligned}\lambda - \lambda_N &= (S_p^\mu\phi, \phi)_2 - (S_{p,N}^\mu\phi_N, \phi_N)_2 \\ &= (S_p^\mu\phi, \phi - \phi_N)_2 + ((S_p^\mu - S_{p,N}^\mu)\phi, \phi_N)_2 + (\phi - \phi_N, S_{p,N}^\mu\phi_N)_2,\end{aligned} \tag{5.15}$$

where the last equality follows from the self-adjointness of $S_{p,N}^\mu$. Therefore we have

$$|\lambda - \lambda_N| \leq (|\lambda| + |\lambda_N|)\,\|\phi - \phi_N\|_2 + \|(S_p^\mu - S_{p,N}^\mu)\phi\|_2. \tag{5.16}$$

Here $\|\phi - \phi_N\|_2$ is bounded by

$$\|\phi - \phi_N\|_2 \leq \|\phi - P_{\lambda_N}(S_{p,N}^\mu)\phi\|_2 + \|P_{\lambda_N}(S_{p,N}^\mu)\phi - \phi_N\|_2. \tag{5.17}$$

Since $\|P_{\lambda_N}(S_{p,N}^\mu)\phi - \phi_N\|_2$ in (5.17) is bounded as

$$\begin{aligned}\|P_{\lambda_N}(S_{p,N}^\mu)\phi - \phi_N\|_2 &= \left\|\left(1 - \frac{1}{\|P_{\lambda_N}(S_{p,N}^\mu)\phi\|_2}\right)P_{\lambda_N}(S_{p,N}^\mu)\phi\right\|_2 \\ &= \Big|\,\|P_{\lambda_N}(S_{p,N}^\mu)\phi\|_2 - 1\,\Big| \\ &\leq \|P_{\lambda_N}(S_{p,N}^\mu)\phi - \phi\|_2,\end{aligned}$$

we have

$$\|\phi - \phi_N\|_2 \leq 2\|\phi - P_{\lambda_N}(S_{p,N}^\mu)\phi\|_2. \tag{5.18}$$

Applying the inequality (5.5) of Lemma 5.1 to (5.18), we have

$$\|\phi - \phi_N\|_2 \leq \frac{4}{r_\lambda}\|(S_p^\mu - S_{p,N}^\mu)\phi\|_2. \tag{5.19}$$

Then combining (5.16) and (5.19), we have

$$|\lambda - \lambda_N| \leq \left(\frac{4}{r_\lambda}(|\lambda| + |\lambda_N|) + 1\right)\|(S_p^\mu - S_{p,N}^\mu)\phi\|_2.$$

Finally, noting $|\lambda_N| \leq |\lambda - \lambda_N| + |\lambda| \leq r_\lambda/2 + |\lambda|$, we have the conclusion. ☐

Remark 5.1. *Assumption 1, i.e. the self-adjointness of S_p^μ is used (i) to estimate the norm of the resolvent of the approximate operator as $\|(\nu I - S_{p,N}^\mu)^{-1}\|_2 = \mathrm{dist}(\nu, \sigma(S_{p,N}^\mu))^{-1}$ in (5.8), (ii) to make an approximate eigenvector as (5.14) and (iii) to deduce the equality (5.15).*

What remains is to estimate $\|(S_p^\mu - S_{p,N}^\mu)\phi\|_2$ in the RHS of (5.13). Since

$$\begin{aligned}\|(S_p^\mu - S_{p,N}^\mu)\phi\|_2 &= \|(S_p^\mu - \hat{P}_N S_p^\mu)\phi + (\hat{P}_N S_p^\mu - \hat{P}_N S_p^\mu \hat{P}_N)\phi\|_2 \\ &\leq \|(I - \hat{P}_N) S_p^\mu \phi\|_2 + \|\hat{P}_N S_p^\mu (I - \hat{P}_N)\phi\|_2 \\ &\leq (|\lambda| + \|\hat{P}_N S_p^\mu\|_2)\|(I - \hat{P}_N)\phi\|_2,\end{aligned}$$

we consider estimation of $\|\hat{P}_N S_p^\mu\|_2$ and $\|(I-\hat{P}_N)\phi\|_2$. Their estimates are given by the following lemmas, whose proofs are shown in Appendix A. It suffices to consider the case of Assumption 2a for $\|\hat{P}_N S_p^\mu\|_2$, whereas the both cases of Assumptions 2a and 2b need to be considered for $\|(I - \hat{P}_N)\phi\|_2$ which depends on the smoothness of ϕ.

Lemma 5.3. *Let S_p^μ satisfy Assumption 2a. Then there exits a positive real constant C depending only on p and $f_0, f_1, \ldots, f_{p-1}$ such that*

$$\|\hat{P}_N S_p^\mu\|_2 \leq C\, N^{p+1/2}.$$

Lemma 5.4. *Let S_p^μ satisfy Assumption 2a and $\phi \in H^p([0,L])_{\mathrm{per}}$ be an eigenvector of S_p^μ corresponding to $\lambda \in \sigma(S_p^\mu)$ with $\|\phi\|_2 = 1$. Then for any positive integer q there exists a positive constant C depending only on ϕ and q such that*

$$\|(I - \hat{P}_N)\phi\|_2 \leq C\,(1+N)^{-q}.$$

Lemma 5.5. *Let S_p^μ satisfy Assumption 2b for $d > 0$ and $\phi \in H^p([0,L])_{\mathrm{per}}$ be an eigenvector of S_p^μ corresponding to $\lambda \in \sigma(S_p^\mu)$ with $\|\phi\|_2 = 1$. Then for any ε with $0 < \varepsilon < d$ there exists a positive constant C depending only on ϕ, d and ε such that*

$$\|(I - \hat{P}_N)\phi\|_2 \leq C\, \exp\left(-\frac{2\pi\,(d-\varepsilon)}{L}\, N\right),$$

where L is the period of the coefficient functions of S_p^μ.

Here we prove Theorem 3.4.

Proof of Theorem 3.4. Since it follows from Lemmas 5.3 and 5.4 that $\|(S_p^\mu - S_{p,N}^\mu)\phi\|_2 \to 0$ as $N \to \infty$, the condition (5.12) in Lemma 5.2 is satisfied. Therefore we can deduce the conclusion (3.8) from Lemmas 5.2, 5.3 and 5.4, and (3.9) from Lemmas 5.2, 5.3 and 5.5. ☐

5.2 Proof of Theorem 3.5

We use almost the same methods as the ones of Lemmas 5.1 and 5.2. On Assumption 3b, we consider $P_{\Lambda_N}(S_{p,N}^\mu)\phi$ for an eigenvector ϕ corresponding to $\lambda \in \sigma(S_{p,N}^\mu)$, where $\Lambda_N = \{\lambda_{N,1}, \ldots, \lambda_{N,k}\}$,

$$P_{\Lambda_N}(S_{p,N}^\mu) = \frac{1}{2\pi \mathrm{i}} \oint_{\Gamma_\Lambda(S_p^\mu)} R_\mu(S_{p,N}^\mu)\, \mathrm{d}\mu, \tag{5.20}$$

and $\Gamma_\Lambda(S_p^\mu)$ is a directed circle with center λ and radius r_λ, which contains Λ_N. Here it should be noted that (5.2), (5.3) and (5.4) can be used for Assumption 3b when λ_N is replaced by $\lambda_{N,i}$ $(i = 1, 2, \ldots, k)$. In this case $P_{\Lambda_N}(S_{p,N}^\mu)\phi$ is not an eigenvector corresponding to a single eigenvalue of $S_{p,N}^\mu$ but a linear combination of eigenvectors corresponding to $\lambda_{N,1}, \ldots, \lambda_{N,k}$. Then we can estimate the convergence rate of one of them after appropriate rearrangement of them explained later.

Lemma 5.6. *Let S_p^μ satisfy Assumptions 1 and 2a, $\lambda \in \sigma(S_p^\mu)$ and $\lambda_{N,1}, \ldots, \lambda_{N,k} \in \sigma(S_{p,N}^\mu)$ satisfy Assumption 3b. Then for an eigenvector ϕ corresponding to λ we have*

$$\|\phi - P_{\Lambda_N}(S_{p,N}^\mu)\phi\|_2 \leq \frac{2}{r_\lambda}\|(S_p^\mu - S_{p,N}^\mu)\phi\|_2, \tag{5.21}$$

where $\varLambda_N = \{\lambda_{N,1}, \dots, \lambda_{N,k}\}$. Furthermore, if $\|\phi\|_2 = 1$ and there exists a positive integer N_0 such that for any $N > N_0$

$$\frac{2}{r_\lambda}\|(S_p^\mu - S_{p,N}^\mu)\phi\|_2 < 1, \tag{5.22}$$

we have $P_{\varLambda_N}(S_{p,N}^\mu)\phi \neq 0$ for $N > N_0$.

Proof. Noting the contour $\varGamma_\lambda(S_p^\mu) = \partial B_\lambda(r_\lambda)$ surrounds λ and $\lambda_{N,1}, \dots, \lambda_{N,k}$ in the case of Assumption 3b, we can use $\varGamma_\lambda(S_p^\mu)$ to define $P_\lambda(S_p^\mu)$ and $P_{\varLambda_N}(S_{p,N}^\mu)$. Then we can obtain the conclusions in a similar manner to Lemma 5.1. □

Lemma 5.7. *Let S_p^μ satisfy Assumptions 1 and 2a, $\lambda \in \sigma(S_p^\mu)$ and $\lambda_{N,1}, \dots, \lambda_{N,k} \in \sigma(S_{p,N}^\mu)$ satisfy Assumption 3b. Furthermore, for an eigenvector ϕ corresponding to λ with $\|\phi\|_2 = 1$, assume that there exists a positive integer N_0 such that for any $N > N_0$*

$$\frac{2}{r_\lambda}\|(S_p^\mu - S_{p,N}^\mu)\phi\|_2 < 1. \tag{5.23}$$

Then for $N > N_0$ there exist a choice of an index $i_N \in \{1, \dots, k\}$ and an integer k'_N with $2 \leq k'_N \leq k$ such that

$$|\lambda - \lambda_{N,i_N}| \leq \left\{K_{\lambda,N}\,\|(S_p^\mu - S_{p,N}^\mu)\phi\|_2\right\}^{1/k'_N}, \tag{5.24}$$

where

$$\begin{aligned} K_{\lambda,N} = (4|\lambda| + r_\lambda)\Bigg[&k^2 \max\left\{1, \left(|\lambda| + \frac{r_\lambda}{2}\right)^{2k}\right\} \max\left\{1, \|S_{p,N}^\mu\|_2^k\right\} \\ &+ \frac{2}{r_\lambda}\max\left\{1, \left(\|S_{p,N}^\mu\|_2 + |\lambda| + \frac{r_\lambda}{2}\right)^k\right\}\Bigg] + \max\left\{1, \left(\frac{r_\lambda}{2}\right)^k\right\}. \end{aligned}$$

In particular, if there exists $N_1 \geq N_0$ such that $K_{\lambda,N}\,\|(S_p^\mu - S_{p,N}^\mu)\phi\|_2 < 1$ for $N > N_1$, the integer k'_N in (5.24) can be replaced by k.

Proof. If there exist $i, j \in \{1, \dots, k\}$ with $i \neq j$ such that $\lambda_{N,i} = \lambda_{N,j}$, we reserve only one of them, and consequently we obtain the set of indices Ω_N such that $\{\lambda_{N,i} \mid i \in \Omega_N\}$ is identical to $\{\lambda_{N,1}, \dots, \lambda_{N,k}\}$ as a set and the elements of it are mutually distinct. Then we renumber the elements of $\{\lambda_{N,i} \mid i \in \Omega_N\}$ and let it be denoted by $\{\lambda_{N,1}, \dots, \lambda_{N,k'_N}\}$, where $k'_N = \#\Omega_N$. Without loss of generality we can assume that $2 \leq k'_N$.

It follows from Lemma 5.6 that $P_{\varLambda_N}(S_{p,N}^\mu)\phi$ is a nonzero linear combination of eigenvectors $\phi_{N,1}, \dots, \phi_{N,k'_N}$ corresponding to $\lambda_{N,1}, \dots, \lambda_{N,k'_N}$ for $N > N_0$. Then there exists a sequence of coefficients $a_{N,1}, \dots, a_{N,k'_N} \in \mathbf{C}$ with $(a_{N,1}, \dots, a_{N,k'_N}) \neq (0, \dots, 0)$ such that

$$P_{\varLambda_N}(S_{p,N}^\mu)\phi = a_{N,1}\phi_{N,1} + \cdots + a_{N,k'_N}\phi_{N,k'_N} \tag{5.25}$$

for $N > N_0$. Here we define $i_N(1)$ as a minimizer of $|\lambda - \lambda_{N,i}|$ for i with $a_{N,i} \neq 0$ in (5.25), i.e.

$$|\lambda - \lambda_{N,i_N(1)}| = \min\{|\lambda - \lambda_{N,i}| \mid i : a_{N,i} \neq 0 \text{ in (5.25)}\}. \tag{5.26}$$

Moreover, we define $\{i_N(2), \dots, i_N(k'_N)\}$ as a permutation of $\{1, \dots, k'_N\} \setminus \{i_N(1)\}$. Consequently, we obtain the ordered set $\{\lambda_{N,i_N(1)}, \dots, \lambda_{N,i_N(k'_N)}\}$ for $N > N_0$. For simplicity, let $\lambda_{N,j}$, $\phi_{N,j}$ and $a_{N,j}$ denote $\lambda_{N,i_N(j)}$, $\phi_{N,i_N(j)}$ and $a_{N,i_N(j)}$ for $j = 1, \dots, k'_N$, respectively. In the following we consider (5.25) under this renumbering.

Here we begin the estimate of the convergence rate of $\lambda_{N,1}$. Applying the operator $(S_{p,N}^\mu - \lambda_{N,2}I)\cdots(S_{p,N}^\mu - \lambda_{N,k'_N}I)$ to the both side of (5.25), we have

$$g_N(S_{p,N}^\mu)\,P_{\varLambda_N}(S_{p,N}^\mu)\,\phi = a_{N,1}\,g_N(\lambda_{N,1})\,\phi_{N,1},$$

where g_N is the polynomial defined as

$$g_N(x) = (x - \lambda_{N,2})\cdots(x - \lambda_{N,k'_N}). \tag{5.27}$$

Because $a_{N,1} \neq 0$ and $\lambda_{N,1} \neq \lambda_{N,i}$ $(i = 2, \ldots, k'_N)$, we can define a normalized eigenvector ϕ_N corresponding to $\lambda_{N,1}$ as

$$\phi_N = C_N^{-1}\, g_N(S^{\mu}_{p,N})\, P_{\Lambda_N}(S^{\mu}_{p,N})\, \phi, \tag{5.28}$$

where

$$C_N = \mathrm{sign}(g_N(\lambda))\, \|\, g_N(S^{\mu}_{p,N})\, P_{\Lambda_N}(S^{\mu}_{p,N})\, \phi\|_2, \tag{5.29}$$

Using ϕ_N in (5.28), in the same manner as (5.16), we can derive an estimate:

$$|\lambda - \lambda_{N,1}| \leq (|\lambda| + |\lambda_{N,1}|)\|\phi - \phi_N\|_2 + \|(S^{\mu}_p - S^{\mu}_{p,N})\phi\|_2. \tag{5.30}$$

Multiplying the both side of (5.30) by $|g_N(\lambda)|$, we have

$$\prod_{i=1}^{k'_N} |\lambda - \lambda_{N,i}| \leq (|\lambda| + |\lambda_{N,1}|)\, |g_N(\lambda)|\, \|\phi - \phi_N\|_2 + |g_N(\lambda)|\, \|(S^{\mu}_p - S^{\mu}_{p,N})\phi\|_2. \tag{5.31}$$

Noting Assumption 3b and (5.26), we can deduce from (5.31) that

$$|\lambda - \lambda_{N,1}|^{k'_N} \leq (|\lambda| + |\lambda_{N,1}|)\, |g_N(\lambda)|\, \|\phi - \phi_N\|_2 + (r_\lambda/2)^{k'_N - 1}\|(S^{\mu}_p - S^{\mu}_{p,N})\phi\|_2. \tag{5.32}$$

Then what remains is to estimate $|g_N(\lambda)|\, \|\phi - \phi_N\|_2$. By the triangle inequality, we have

$$|g_N(\lambda)|\, \|\phi - \phi_N\|_2 \leq \|g_N(\lambda)\phi - C_N\phi_N\|_2 + \|(C_N - g_N(\lambda))\, \phi_N\|_2\,,$$

and

$$\begin{aligned}
&\|g_N(\lambda)\phi - C_N\phi_N\|_2 \\
&= \|g_N(S^{\mu}_p)\, P_\lambda(S^{\mu}_p)\, \phi - g_N(S^{\mu}_{p,N})\, P_{\Lambda_N}(S^{\mu}_{p,N})\, \phi\|_2 \\
&\leq \|\{g_N(S^{\mu}_p) - g_N(S^{\mu}_{p,N})\}\, \phi\|_2 + \|g_N(S^{\mu}_{p,N})\{P_\lambda(S^{\mu}_p) - P_{\Lambda_N}(S^{\mu}_{p,N})\}\, \phi\|_2.
\end{aligned} \tag{5.33}$$

Therefore setting

$$\begin{aligned}
E_{1,N} &= \|\, (C_N - g_N(\lambda))\, \phi_N\|_2, \\
E_{2,N} &= \|\{g_N(S^{\mu}_p) - g_N(S^{\mu}_{p,N})\}\, \phi\|_2, \\
E_{3,N} &= \|g_N(S^{\mu}_{p,N})\{P_\lambda(S^{\mu}_p) - P_{\Lambda_N}(S^{\mu}_{p,N})\}\, \phi\|_2
\end{aligned}$$

we have

$$|g_N(\lambda)|\, \|\phi - \phi_N\|_2 \leq E_{1,N} + E_{2,N} + E_{3,N}. \tag{5.34}$$

We can estimate $E_{1,N}$, $E_{2,N}$ and $E_{3,N}$ as follows. Using (5.28) and (5.29), we have

$$\begin{aligned}
E_{1,N} &= |C_N - g_N(\lambda)| = |C_N - \mathrm{sign}(g_N(\lambda))\|g_N(\lambda)\phi\|_2| \\
&= |\|C_N\phi_N\|_2 - \|g_N(\lambda)\phi\|_2| \leq \|C_N\phi_N - g_N(\lambda)\phi\|_2 \\
&\leq E_{2,N} + E_{3,N},
\end{aligned} \tag{5.35}$$

where the last inequality is due to (5.33). Noting that ϕ is an eigenvector of S^{μ}_p corresponding to λ, we have

$$\begin{aligned}
\{g_N(S^{\mu}_p) - g_N(S^{\mu}_{p,N})\}\phi &= \{g_N(\lambda) - g_N(S^{\mu}_{p,N})\}\phi \\
&= \sum_{i=0}^{k'_N - 1} \alpha(i, \Lambda_N)\left(\lambda^{k'_N - 1 - i} I - (S^{\mu}_{p,N})^{k'_N - 1 - i}\right)\phi \\
&= \sum_{i=0}^{k'_N - 2} \alpha(i, \Lambda_N)\, \beta(i, \lambda, S^{\mu}_{p,N})\, (\lambda I - S^{\mu}_{p,N})\, \phi \\
&= \sum_{i=0}^{k'_N - 2} \alpha(i, \Lambda_N)\, \beta(i, \lambda, S^{\mu}_{p,N})\, (S^{\mu}_p - S^{\mu}_{p,N})\, \phi,
\end{aligned} \tag{5.36}$$

where $\alpha(i,\Lambda_N) = (-1)^i \sum_{2\le j_1<\cdots<j_i\le k'_N} \lambda_{N,j_1}\cdots\lambda_{N,j_i}$ and $\beta(i,\lambda,S^\mu_{p,N}) = \sum_{l=0}^{k'_N-2-i} \lambda^{k'_N-2-i-l}(S^\mu_{p,N})^l$. Therefore noting

$$|\alpha(i,\Lambda_N)| \le \left(|\lambda| + \frac{r_\lambda}{2}\right)^i \le \max\left\{1, \left(|\lambda| + \frac{r_\lambda}{2}\right)^{k'_N}\right\},$$

$$\begin{aligned}\|\beta(i,\lambda,S^\mu_{p,N})\|_2 &\le \max\left\{1, |\lambda|^{k'_N}\right\} \sum_{l=0}^{k'_N-2-i} \|S^\mu_{p,N}\|_2^l \\ &\le \max\left\{1, |\lambda|^{k'_N}\right\} k'_N \max\left\{1, \|S^\mu_{p,N}\|_2^{k'_N}\right\}\end{aligned}$$

and (5.36), we have

$$E_{2,N} \le (k'_N)^2 \max\left\{1, \left(|\lambda| + \frac{r_\lambda}{2}\right)^{2k'_N}\right\} \max\left\{1, \|S^\mu_{p,N}\|_2^{k'_N}\right\} \|(S^\mu_p - S^\mu_{p,N})\phi\|_2. \tag{5.37}$$

Using (5.21) in Lemma 5.6, we have

$$\begin{aligned}E_{3,N} &\le \left(\prod_{i=2}^{k'_N} \|S^\mu_{p,N} - \lambda_{N,i} I\|_2\right) \|\{P_\lambda(S^\mu_p) - P_{\Lambda_N}(S^\mu_{p,N})\}\phi\|_2 \\ &\le \left(\prod_{i=2}^{k'_N} (\|S^\mu_{p,N}\|_2 + |\lambda_{N,i}|)\right) \|\phi - P_{\Lambda_N}(S^\mu_{p,N})\phi\|_2 \\ &\le \left(\|S^\mu_{p,N}\|_2 + |\lambda| + \frac{r_\lambda}{2}\right)^{k'_N-1} \frac{2}{r_\lambda} \|(S^\mu_p - S^\mu_{p,N})\phi\|_2. \end{aligned} \tag{5.38}$$

Finally, noting $k'_N \le k$ and combining (5.32), (5.34), (5.35), (5.37) and (5.38), we have

$$\begin{aligned}&|\lambda - \lambda_{N,1}|^{k'_N} \\ &\le 2(|\lambda| + |\lambda_{N,1}|)(E_{2,N} + E_{3,N}) + (r_\lambda/2)^{k'_N-1} \|(S^\mu_p - S^\mu_{p,N})\phi\|_2. \\ &\le \left[(4|\lambda| + r_\lambda)\left[k^2 \max\left\{1, \left(|\lambda| + \frac{r_\lambda}{2}\right)^{2k}\right\} \max\left\{1, \|S^\mu_{p,N}\|_2^k\right\}\right.\right. \\ &\quad \left.\left. + \frac{2}{r_\lambda} \max\left\{1, \left(\|S^\mu_{p,N}\|_2 + |\lambda| + \frac{r_\lambda}{2}\right)^k\right\}\right] + \max\left\{1, \left(\frac{r_\lambda}{2}\right)^k\right\}\right] \|(S^\mu_p - S^\mu_{p,N})\phi\|_2.\end{aligned}$$

Thus we obtain the conclusion. ☐

Here we prove Theorem 3.5.

Proof of Theorem 3.5. In a similar manner to the proof of Theorem 3.4, we can deduce the conclusion (3.10) from Lemmas 5.3, 5.4 and 5.7, and (3.11) from Lemmas 5.3, 5.5, and 5.7. Note that $\|S^\mu_{p,N}\|_2 \le \|\hat{P}_N S^\mu_p\|_2 \|\hat{P}_N\|_2 = \|\hat{P}_N S^\mu_p\|_2$. ☐

5.3 Proof of Theorem 3.6

To prove the theorem, we use a similar technique to Lemmas 5.1 and 5.2, in which we exchange the roles of ϕ and ϕ_N. That is, we take an eigenvector ϕ_N with $\|\phi_N\|_2 = 1$ corresponding to λ_N, and using ϕ_N and the projection $P_\lambda(S^\mu_p)$ for $\lambda \in \sigma(S^\mu_p)$, we obtain an eigenvector ϕ with $\|\phi\|_2 = 1$ corresponding to λ. First, we present a lemma guaranteeing the feasibility of such procedure, which corresponds to Lemma 5.1. We omit its proof since it is proved in almost the same manner as Lemma 5.1.

Lemma 5.8. *Let S^μ_p satisfy Assumptions 1 and 2a, $\lambda \in \sigma(S^\mu_p)$ and $\lambda_N \in \sigma(S^\mu_{p,N})$ satisfy Assumption 3a. Then for an eigenvector ϕ_N corresponding to λ_N we have*

$$\|\phi_N - P_\lambda(S^\mu_p)\phi_N\|_2 \le \frac{2}{r_\lambda} \|(S^\mu_p - S^\mu_{p,N})\phi_N\|_2. \tag{5.39}$$

Furthermore, if $\|\phi_N\|_2 = 1$ *and*

$$\frac{2}{r_\lambda}\|(S_p^\mu - S_{p,N}^\mu)\phi_N\|_2 < 1, \tag{5.40}$$

we have $P_\lambda(S_p^\mu)\phi_N \neq 0$.

Next, we present a counterpart of Lemma 5.2, whose proof is also omitted.

Lemma 5.9. *Let* S_p^μ *satisfy Assumptions 1 and 2a,* $\lambda \in \sigma(S_p^\mu)$ *and* $\lambda_N \in \sigma(S_{p,N}^\mu)$ *satisfy Assumption 3a. Furthermore, for an eigenvector* ϕ_N *corresponding to* λ_N *with* $\|\phi_N\|_2 = 1$*, assume that*

$$\frac{2}{r_\lambda}\|(S_p^\mu - S_{p,N}^\mu)\phi_N\|_2 < 1. \tag{5.41}$$

Then we have

$$|\lambda - \lambda_N| \leq \left(3 + \frac{8|\lambda|}{r_\lambda}\right)\|(S_p^\mu - S_{p,N}^\mu)\phi_N\|_2. \tag{5.42}$$

Here we prove Theorem 3.6.

Proof of Theorem 3.6. It follows from the assumption (3.12), Theorems 3.1 and 3.2 that the set of the accumulation points of the sequence $\{\lambda_N\}$ coincides with $\sigma(S_p^\mu) \cap B_\zeta(r)$. Moreover, the elements of $\sigma(S_p^\mu) \setminus B_\zeta(r)$ do not exist in the interior of $B_\zeta(9r)$. In the following we set $\Lambda_{\zeta,r} = \sigma(S_p^\mu) \cap B_\zeta(r)$.

First, we show that $\{\lambda_N\}$ is convergent and $\lambda \in \Lambda_{\zeta,r}$ is uniquely determined as the limit of $\{\lambda_N\}$. To prove $\{\lambda_N\}$ is Cauchy, we choose an arbitrary integer $M > 0$ and estimate $|\lambda_{N+M} - \lambda_N|$ for sufficiently large N. For an eigenvalue $\lambda \in \Lambda_{\zeta,r}$, we take an eigenvector ϕ with $\|\phi\|_2 = 1$ corresponding to λ. Then ϕ satisfies

$$P_{\Lambda_{\zeta,r}}(S_p^\mu)\phi = \frac{1}{2\pi \mathrm{i}}\oint_{\Gamma_{\zeta,r}} (\nu I - S_p^\mu)^{-1}\phi \,\mathrm{d}\nu = \frac{1}{2\pi \mathrm{i}}\oint_{\Gamma_{\zeta,r}} (\nu - \lambda)^{-1}\phi \,\mathrm{d}\nu = \phi, \tag{5.43}$$

where $\Gamma_{\zeta,r}$ is the boundary of $B_\zeta(2r)$ with counterclockwise direction. Note that (5.43) holds true regardless of the number of the elements in $\Lambda_{\zeta,r}$. Using this ϕ, we define eigenvectors corresponding to λ_{N+M} and λ_N as

$$\phi_{N+M} = \frac{1}{\|P_{\lambda_{N+M}}(S_{p,N+M}^\mu)\phi\|_2} P_{\lambda_{N+M}}(S_{p,N+M}^\mu)\phi$$

and

$$\phi_N = \frac{1}{\|P_{\lambda_N}(S_{p,N}^\mu)\phi\|_2} P_{\lambda_N}(S_{p,N}^\mu)\phi,$$

respectively. Here we use again $\Gamma_{\zeta,r}$ to define $P_{\lambda_{N+M}}(S_{p,N+M}^\mu)$ and $P_{\lambda_N}(S_{p,N}^\mu)$. The availability of the eigenvectors can be guaranteed by a similar manner to Lemma 5.1. Then noting $S_{p,N}^\mu = \hat{P}_N S_p^\mu \hat{P}_N$ etc. we have

$$\begin{aligned}
\lambda_{N+M} - \lambda_N &= (S_{p,N+M}^\mu \phi_{N+M}, \phi_{N+M}) - (S_{p,N+M}^\mu \phi_N, \phi_N) \\
&= (S_{p,N+M}^\mu \phi_{N+M}, \phi_{N+M} - \phi_N) + (S_{p,N+M}^\mu(\phi_{N+M} - \phi_N), \phi_N) \\
&= (S_{p,N+M}^\mu \phi_{N+M}, \phi_{N+M} - \phi_N) + (\phi_{N+M} - \phi_N, S_{p,N+M}^\mu \phi_N)
\end{aligned}$$

and therefore

$$|\lambda_{N+M} - \lambda_N| \leq 2\|S_{p,N+M}^\mu\|_2(\|\phi_{N+M} - \phi\|_2 + \|\phi - \phi_N\|_2). \tag{5.44}$$

Using a similar manner to Lemma 5.2, Lemma 5.3 and Lemma 5.4, we can show that the RHS of (5.44) is $\mathrm{O}(N^{-q})$ for any positive integer q. In fact, $\|S^{\mu}_{p,N+M}\|_2 = \mathrm{O}(N^{p+1/2})$ by Lemma 5.3 and

$$\begin{aligned}\|\phi - \phi_N\|_2 &\leq 2\,\|\phi - P_{\lambda_N}(S^{\mu}_{p,N})\phi\|_2 = 2\,\|P_{\Lambda_{\zeta,r}}(S^{\mu}_p)\phi - P_{\lambda_N}(S^{\mu}_{p,N})\phi\|_2 \\ &\leq 4r \max_{\nu\in\Gamma_{\zeta,r}} \|(\nu I - S^{\mu}_{p,N})^{-1}(S^{\mu}_{p,N} - S^{\mu}_p)(\nu I - S^{\mu}_p)^{-1}\phi\|_2 \\ &= 4r \max_{\nu\in\Gamma_{\zeta,r}} \|(\nu I - S^{\mu}_{p,N})^{-1}(S^{\mu}_{p,N} - S^{\mu}_p)(\nu - \lambda)^{-1}\phi\|_2 \\ &\leq 4r \max_{\nu\in\Gamma_{\zeta,r}} \frac{1}{|\nu-\lambda|}\frac{1}{|\nu-\lambda_N|}\|(S^{\mu}_{p,N} - S^{\mu}_p)\phi\|_2 \\ &\leq \frac{4}{r}\|(S^{\mu}_{p,N} - S^{\mu}_p)\phi\|_2.\end{aligned}$$

Thus we can show that $|\lambda_{N+M} - \lambda_N| \to 0$ as $N \to \infty$ and $\lambda \in \Lambda_{\zeta,r}$ is uniquely determined. Combining this fact and $|\lambda - \lambda_N| \leq 2r$, we can show that Assumption 3a is satisfied for $\{\lambda_N\}$, λ, and $r_\lambda = 4r$. In fact, for any $\tilde{\lambda} \in \sigma(S^{\mu}_p) \cup \sigma(S^{\mu}_{p,N}) \setminus \{\lambda, \lambda_N\}$ we have

$$|\tilde{\lambda} - \lambda| \geq |\tilde{\lambda} - \zeta| - |\lambda - \zeta| \geq 2r_\lambda.$$

Next, we show the error bound (3.13). By Lemma 5.9, it follows from (5.42) and $|\lambda| \leq r + |\zeta|$ that

$$|\lambda - \lambda_N| \leq \left(5 + \frac{3|\zeta|}{r}\right)\|(S^{\mu}_p - S^{\mu}_{p,N})\phi_N\|_2 \tag{5.45}$$

for $\lambda \in \sigma(S^{\mu}_p) \cap B_\zeta(r)$. Furthermore, we have

$$\begin{aligned}(S^{\mu}_p - S^{\mu}_{p,N})\phi_N &= (I - \hat{P}_N)S^{\mu}_p\phi_N \\ &= \sum_{|l|>N}\left\{\sum_{j=0}^{p}\sum_{m=l-N}^{l+N}(\hat{f}_j)_m(\hat{\phi}_N)_{l-m}\frac{1}{\sqrt{L}}\left(-\mathrm{i}\frac{2\pi(l-m)}{L}\right)^j\right\}\frac{1}{\sqrt{L}}\exp\left(-\mathrm{i}\frac{2\pi l}{L}x\right),\end{aligned}$$

and therefore

$$\begin{aligned}\|(S^{\mu}_p - S^{\mu}_{p,N})\phi_N\|_2 &\leq \sum_{|l|>N}\left|\sum_{j=0}^{p}\sum_{m=l-N}^{l+N}(\hat{f}_j)_m(\hat{\phi}_N)_{l-m}\frac{1}{\sqrt{L}}\left(-\mathrm{i}\frac{2\pi(l-m)}{L}\right)^j\right| \\ &\leq \frac{1}{\sqrt{L}}\left(\frac{2\pi N}{L}\right)^p\sum_{j=0}^{p}\sum_{|l|>N}\sum_{m=l-N}^{l+N}\left|(\hat{f}_j)_m(\hat{\phi}_N)_{l-m}\right|.\end{aligned} \tag{5.46}$$

Considering the case $|l| \geq 2N$, we have

$$\sum_{|l|\geq 2N}\sum_{m=l-N}^{l+N}\left|(\hat{f}_j)_m(\hat{\phi}_N)_{l-m}\right| \leq \sum_{|l|\geq 2N}\sum_{m=l-N}^{l+N}\left|(\hat{f}_j)_m\right| \leq (2N+1)\sum_{|m|\geq N}\left|(\hat{f}_j)_m\right|. \tag{5.47}$$

Combining (5.45), (5.46) and (5.47), we obtain the error bound (3.13). □

6 Concluding Remarks

We have considered the convergence rates and the explicit error bounds of Hill's method, which is a numerical method for computing the spectra of self-adjoint differential operators S_p in (1.1) with periodic coefficient functions $\tilde{f}_j$. On the assumption (Assumptions 3a and 3b) that the computed eigenvalue λ_N is close to the exact one λ, it is shown in Theorems 3.4 and 3.5 that the convergence rate of the computed eigenvalue is all order polynomial in the case of $\tilde{f}_j$ being C^∞ as shown in (3.8) and (3.10), and exponential in the case of $\tilde{f}_j$ being analytic as shown in (3.9) and (3.11). In addition, even if the exact eigenvalue λ is unknown, it is shown in Theorems 3.4, 3.5 and 3.6 that, if the condition (3.12) is satisfied, we can obtain the convergence rate using (3.8)–(3.11) and the explicit error bound using (3.13). There are some cases in which the condition (3.12) can be checked using Gershgorin's theorem. These theorems are proved using the Dunford integrals $P_\lambda(S^{\mu}_{p,N})$ in (5.1) and $P_\lambda(S^{\mu}_{p})$ in the proof of Theorem 3.6, which project an eigenvector to the corresponding eigenspace. This integral is suitable for proving these theorems, because the boundary of the neighborhood of an eigenvalue can be directly used as the contour of the integral. Numerical examples using Hill's operator (4.1) support these theoretical results.

As described in Remark 5.1, self-adjointness of the operator S^{μ}_{p} enables us to estimate the norm of the resolvent as shown in (5.8), and to project an eigenvector to the corresponding eigenspace using the Dunford integral. For the case of non-self-adjoint operators, we have developed some other approaches, which will be reported somewhere else soon.

Acknowledgments

This work is partially supported by Grant-in-Aid for Scientific Research (S) 20224001.

References

[1] Atkinson. K., The numerical solution of the eigenvalue problem for compact integral operators, Trans. Amer. Math. Soc. **129**, 458–465 (1967)

[2] Atkinson, K., Convergence rates for approximate eigenvalues of compact integral operators, SIAM J. Num. Anal. **12**, 213–222 (1975)

[3] Atkinson, K., Han, W., Theoretical numerical analysis: A functional analysis framework, second edition, Springer, New York (2005)

[4] Curtis, C. W., Deconinck, B., On the convergence of Hill's method, Math. Comp. **79**, 169–187 (2010)

[5] Dunford, N, Schwartz, J. T., Linear operators Part II Spectral theory, Self adjoint operators in Hilbert space, Wiley & Sons, New Jersey (1963)

[6] Dunford, N, Schwartz, J. T., Linear operators Part III Spectral operators, Wiley & Sons, New Jersey (1971)

[7] Deconinck, B., Kutz, J. N., Computing spectra of linear operators using Hill's method, J. Computational Physics **219**, 296–321 (2006)

[8] Hill, G. W., On the part of the motion of the lunar perigee which is a function of the mean motions of the sun and moon, Acta Math. **8**, 1–36 (1886)

[9] Hille, E., Ordinary differential equations in the complex domain, Dover, New York (1997)

[10] Johnson, M. A., Zumbrun, K., Convergence of Hill's method for nonselfadjoint operators, SIAM J. Numer. Anal. **50**, 64–78 (2012)

[11] Krasnosel'skii, M. A., Vainikko, G. M., Zabreiko, P. P., Rutitskii, Ya. B., Stetsenko, V. Ya., Approximate solution of operator equations, Wolters-Noordhoff, Groningen (1972)

[12] Reed, M., Simon, B., Methods of modern mathematical physics. I. Functional analysis, revised and enlarged edidition, Academic Press, New York (1980)

[13] Reed, M., Simon, B., Methods of modern mathematical physics. IV. Analysis of operators, Academic Press, New York (1978)

[14] Vainikko, G. M., A perturbed Galerkin method and the general theory of approximate methods for nonlinear equations, Zh. Vychisl. Mat. Mat. Fiz. **7**, 723–751 (1967) (in Russian), U. S. S. R. Comput. Math. and Math. Phys. **7**, 18–32 (1967) (in English).

[15] Varga, R. S., Gershgorin and his circles, Springer, Berlin Heidelberg (2004)

A Proofs of Lemmas 5.3, 5.4, and 5.5

First, we prove Lemma 5.3.

Proof of Lemma 5.3. We set $f_p \equiv 1$ for simplicity. Let e_m $(m = 0, \pm 1, \pm 2, \ldots)$ be the Fourier bases defined as

$$\mathrm{e}_m(x) = \frac{1}{\sqrt{L}} \exp\left(-\mathrm{i}\frac{2\pi m}{L}x\right),$$

and let $(\hat{f}_j)_m$ $(j = 0, 1, \ldots, p,\ m = 0, \pm 1, \pm 2, \ldots)$ denote the Fourier coefficients of f_j $(j = 0, 1, \ldots, p)$, i.e. $(\hat{f}_j)_m = (f_j, \mathrm{e}_m)_2$. Then it is a standard fact that under Assumption 2a, for $j = 0, 1, \ldots, p$, $m = 0, \pm 1, \pm 2, \ldots$ and any positive integer q, there exists a positive real constant $C_{j,q}$ depending only on f_j and q such that

$$|(\hat{f}_j)_m| \leq C_{j,q}\,(1 + |m|)^{-q}. \tag{A.1}$$

We estimate the norm of

$$(\hat{P}_N S_p^\mu \psi)(x) = \sum_{n=-N}^{N} \sum_{j=0}^{p} \left\{ \sum_{m=-\infty}^{\infty} (\hat{f}_j)_m \left(-\mathrm{i}\frac{2\pi(n-m)}{L}\right)^j \hat{\psi}_{n-m} \right\} \mathrm{e}_n(x) \tag{A.2}$$

for $N \geq 1$ and a nonzero vector $\psi \in L_2([0, L])_{\mathrm{per}}$. Setting

$$c_j(n, m) = (\hat{f}_j)_m \left(-\mathrm{i}\frac{2\pi(n-m)}{L}\right)^j$$

and $q = p + 1$ in (A.1), for $j = 0, 1, \ldots, p$, $m = 0, \pm 1, \pm 2, \ldots$ and n with $|n| \leq N$, we have

$$\begin{aligned} N^{-j}|c_j(n, m)| &= \left(\frac{2\pi}{L}\right)^j |(\hat{f}_j)_m| \left(\frac{|n| + |m|}{N}\right)^j \\ &\leq \left(\frac{2\pi}{L}\right)^j C_{j,p+1}\,(1 + |m|)^{-p-1} \left(1 + \frac{|m|}{N}\right)^j \\ &\leq \left(\frac{2\pi}{L}\right)^j C_{j,p+1}\,(1 + |m|)^{j-p-1}, \end{aligned}$$

and therefore

$$|c_j(n, m)| \leq \left(\frac{2\pi}{L}\right)^j \frac{C_{j,p+1}}{1 + |m|} N^j. \tag{A.3}$$

Combining (A.2) and (A.3) we have

$$\begin{aligned} \|\hat{P}_N S_p^\mu \psi\|_2 &\leq \sum_{j=0}^{p} \left\| \sum_{n=-N}^{N} \left(\sum_{m=-\infty}^{\infty} c_j(n, m)\hat{\psi}_{n-m} \right) \mathrm{e}_n \right\|_2 \\ &= \sum_{j=0}^{p} \left\{ \sum_{n=-N}^{N} \left| \sum_{m=-\infty}^{\infty} c_j(n, m)\hat{\psi}_{n-m} \right|^2 \right\}^{1/2} \\ &\leq \sum_{j=0}^{p} \left\{ \sum_{n=-N}^{N} \left(\sum_{m=-\infty}^{\infty} |c_j(n, m)|^2 \right) \left(\sum_{m=-\infty}^{\infty} |\hat{\psi}_{n-m}|^2 \right) \right\}^{1/2} \\ &\leq \sum_{j=0}^{p} \left\{ \sum_{n=-N}^{N} \left(\frac{2\pi}{L}\right)^{2j} C_{j,p+1}^2 N^{2j} \left(\sum_{m=-\infty}^{\infty} \frac{1}{(1 + |m|)^2} \right) \|\psi\|_2^2 \right\}^{1/2} \\ &\leq \sum_{j=0}^{p} \tilde{C}_{j,p+1}(2N + 1)^{1/2} N^j \|\psi\|_2, \end{aligned} \tag{A.4}$$

where $\tilde{C}_{j,p+1}$ $(j = 0, 1, \dots, p)$ are bounded constants defined as

$$\tilde{C}_{j,p+1} = \left(\frac{2\pi}{L}\right)^j C_{j,p+1} \left(\sum_{m=-\infty}^{\infty} \frac{1}{(1+|m|)^2}\right)^{1/2}.$$

Hence it follows from (A.4) that

$$\|\hat{P}_N S_p^\mu\|_2 \leq \sum_{j=0}^{p} \tilde{C}_{j,p+1}(2N+1)^{1/2} N^j. \tag{A.5}$$

Thus we have the conclusion. ☐

Next, we prove Lemma 5.4 and 5.5 with the aid of the following two lemmas. Since these lemmas are fundamental in the theory of differential equations, we only give the outlines of their proofs in Appendix B.

Lemma A.1. *Let S_p^μ satisfy Assumption 2a and $\phi \in H^p([0,L])_{\mathrm{per}}$ be an eigenvector of S_p^μ corresponding to $\lambda \in \sigma(S_p^\mu)$. Then ϕ is C^∞.*

Lemma A.2. *Let S_p^μ satisfy Assumption 2b for $d > 0$ and $\phi \in H^p([0,L])_{\mathrm{per}}$ be an eigenvector of S_p^μ corresponding to $\lambda \in \sigma(S_p^\mu)$. Then ϕ can be uniquely extended to an analytic function on $\mathcal{D}_d$ with $\phi(z+L) = \phi(z)$ for any $z \in \mathbf{C}$, where $\mathcal{D}_d$ is defined as (3.1) and L is the period of the coefficient functions of S_p^μ.*

Proof of Lemma 5.4. Since ϕ is C^∞ by Lemma A.1, in a similar manner to (A.1), we can use the fundamental fact for the Fourier coefficients that there exists a positive constant $C_{\phi,q}$ depending only on ϕ and q such that

$$|\hat{\phi}_n| \leq C_{\phi,q}\,(1+|n|)^{-q-1}$$

for any integer n. Then we have

$$\begin{aligned}\|(I-\hat{P}_N)\phi\|_2^2 &= \sum_{|n|>N} |\hat{\phi}_n|^2 \leq \sum_{|n|>N} C_{\phi,q}^2\,(1+|n|)^{-2q-2} \\ &\leq \frac{C_{\phi,q}^2}{(1+N)^{2q}} \sum_{|n|>N} \frac{1}{(1+|n|)^2} \leq \frac{C_{\phi,q}^2}{(1+N)^{2q}} \sum_{n=-\infty}^{\infty} \frac{1}{(1+|n|)^2}.\end{aligned}$$

Thus we obtain the conclusion. ☐

Proof of Lemma 5.5. Since ϕ can be regarded as an analytic function on $\mathcal{D}_d$ with $\phi(z+L) = \phi(z)$ by Lemma A.2, there exists a positive constant $C_{\phi,d,\varepsilon}$ depending only on ϕ, d and ε such that

$$|\hat{\phi}_n| \leq C_{\phi,d,\varepsilon} \exp\left(-\frac{2\pi d_\varepsilon}{L}|n|\right) \tag{A.6}$$

for any integer n, where $d_\varepsilon = d - \varepsilon$. In fact, we can show this estimate by the Paley-Wiener type argument. Noting the periodicity $\phi(z+L) = \phi(z)$, by Cauchy's integral theorem we have

$$\begin{aligned}\hat{\phi}_n &= \frac{1}{\sqrt{L}} \int_0^L \phi(x) \exp\left(-\mathrm{i}\frac{2\pi n}{L}x\right) \mathrm{d}x \\ &= \frac{1}{\sqrt{L}} \int_0^L \phi\left(x - \mathrm{sign}(n)\, d_\varepsilon \mathrm{i}\right) \exp\left(-\mathrm{i}\frac{2\pi n}{L}\left(x - \mathrm{sign}(n)\, d_\varepsilon \mathrm{i}\right)\right) \mathrm{d}x \\ &= \exp\left(-\frac{2\pi d_\varepsilon}{L}|n|\right) \frac{1}{\sqrt{L}} \int_0^L \phi\left(x - \mathrm{sign}(n)\, d_\varepsilon \mathrm{i}\right) \exp\left(-\mathrm{i}\frac{2\pi n}{L}x\right) \mathrm{d}x,\end{aligned}$$

where $\mathrm{sign}(n)$ is the sign of n. This expression implies (A.6). Therefore we have

$$\|(I-\hat{P}_N)\phi\|_2^2 = \sum_{|n|>N} |\hat{\phi}_n|^2 \leq \sum_{|n|>N} C_{\phi,d,\varepsilon}^2 \exp(-2\, d_{\varepsilon,L}\, |n|) = \frac{2\, C_{\phi,d,\varepsilon}^2 \exp(-2\, d_{\varepsilon,L}\, (N+1))}{1 - \exp(-2\, d_{\varepsilon,L})},$$

where $d_{\varepsilon,L} = 2\pi d_\varepsilon / L$. Thus we obtain the conclusion. ☐

B Proofs of Lemmas A.1 and A.2

Proof of Lemma A.1. Since ϕ is an eigenvector of S_p^μ corresponding to λ, we have

$$\phi^{(p)}(x) = -\sum_{j=0}^{p-1} f_j(x)\phi^{(j)}(x) + \lambda\phi(x).$$

Then it follows from $\phi \in H^p([0,L])_{\mathrm{per}}$ that $\phi^{(p)} \in H^1([0,L])_{\mathrm{per}}$. Hence $\phi \in H^{p+1}([0,L])_{\mathrm{per}}$. Iterating a similar argument, we have $\phi \in H^q([0,L])_{\mathrm{per}}$ for any positive integer q. Then the conclusion follows from the fundamental embedding relation: $H^q([0,L])_{\mathrm{per}} \hookrightarrow C^{q-1}([0,L])_{\mathrm{per}}$ for any positive integer q, where $C^{q-1}([0,L])_{\mathrm{per}}$ is the set of periodic functions on $[0,L]$ which are $q-1$ times continuously differentiable (see e.g. [3, Proposition 7.5.4]). $\square$

Proof of Lemma A.2. We choose an arbitrary ε with $0 < \varepsilon < d$ and set $d_\varepsilon = d - \varepsilon$. Then $f_0, f_1, \ldots, f_{p-1}$ are analytic on $\mathcal{D}_{d_\varepsilon}$ and continuous on $\overline{\mathcal{D}_{d_\varepsilon}}$. Hence there exists a positive constant M_ε such that $|f_j(z)| \leq M_\varepsilon$ for all $j = 0, 1, \ldots, p-1$ and $z \in \mathcal{D}_{d_\varepsilon}$.

By Lemma A.1, we can assume the existence of an eigenvector $\phi \in C^\infty([0,L])_{\mathrm{per}}$ corresponding to $\lambda \in \sigma(S_p^\mu)$. Then for the fixed λ and each $x_0 \in \mathbf{R}$, we consider the equation $S_p^\mu \psi = \lambda\psi$ as an initial value problem with the condition $\psi^{(n)}(x_0) = \phi^{(n)}(x_0)$ $(n = 0, \ldots, p-1)$. By the matrix $F(z)$ defined as

$$F(z) = \begin{pmatrix} -f_{p-1}(z) & -f_{p-2}(z) & \cdots & -f_1(z) & -(f_0(z) - \lambda) \\ 1 & 0 & \cdots & & 0 \\ 0 & 1 & \cdots & & \vdots \\ \vdots & \ddots & \ddots & & \vdots \\ 0 & \cdots & \cdots & 1 & 0 \end{pmatrix}$$

and the vector $w_0 = (\phi^{(p-1)}(x_0), \phi^{(p-2)}(x_0), \ldots, \phi(x_0))^T$, the initial value problem above is expressed as

$$w'(z) = F(z)w(z), \quad w(x_0) = w_0, \tag{B.1}$$

where $w(z)$ corresponds to $(\psi^{(p-1)}(z), \psi^{(p-2)}(z), \ldots, \psi(z))^T$. The equation (B.1) is equivalently expressed as

$$w(z) = w_0 + \int_{x_0}^{z} F(s)w(s)\mathrm{d}s. \tag{B.2}$$

By the fundamental method of successive approximations [9, Theorem 2.3.1][1], it is shown that the equation (B.1) has a unique analytic solution w on $\mathcal{D}_{d_\varepsilon}$ for each $x_0 \in \mathbf{R}$. Then what remains is to prove the consistency of the solutions for different x_0's and the periodicity of the solution. First, note that for any x_0 the solution w is identical to $v = (\phi^{(p-1)}, \phi^{(p-2)}, \ldots, \phi)^T$ on $\mathbf{R}$. This follows from the condition

$$\|w(x) - v(x)\| = \left\| \int_{x_0}^{x} F(s)(w(s) - v(s))\mathrm{d}s \right\| \leq \sup_{z \in \mathcal{D}_{d_\varepsilon}} \|F(z)\| \int_{x_0}^{x} \|w(s) - v(s)\|\, |\mathrm{d}s|, \tag{B.3}$$

where $\|\cdot\|$ is a norm of $\mathbf{R}^p$. Then if w_1 and w_2 are the solutions for $x_0^{(1)}$ and $x_0^{(2)}$, respectively, we have

$$\begin{aligned} w_1(z) - w_2(z) &= w_1(x_0^{(1)}) - w_2(x_0^{(2)}) + \int_{x_0^{(1)}}^{z} F(s)w_1(s)\mathrm{d}s - \int_{x_0^{(2)}}^{z} F(s)w_2(s)\mathrm{d}s \\ &= w_1(x_0^{(1)}) - w_2(x_0^{(2)}) - \int_{x_0^{(2)}}^{x_0^{(1)}} F(s)w_2(s)\mathrm{d}s + \int_{x_0^{(1)}}^{z} F(s)(w_1(s) - w_2(s))\mathrm{d}s \\ &= v(x_0^{(1)}) - v(x_0^{(2)}) - \int_{x_0^{(2)}}^{x_0^{(1)}} F(s)v(s)\mathrm{d}s + \int_{x_0^{(1)}}^{z} F(s)(w_1(s) - w_2(s))\mathrm{d}s \\ &= \int_{x_0^{(1)}}^{z} F(s)(w_1(s) - w_2(s))\mathrm{d}s \end{aligned}$$

---

[1] In [9, Theorem 2.3.1], the general form $w' = F(z, w)$ is considered. Here we consider the special case that F is linear with respect to w.

for any $z \in \mathcal{D}_{d_\varepsilon}$, which implies $w_1 \equiv w_2$ on $\mathcal{D}_{d_\varepsilon}$ by the same criterion as (B.3). Next, the expressions

$$\begin{aligned} w(z+L) &= v(x_0+L) + \int_{x_0+L}^{z+L} F(s)w(s)\mathrm{d}s = v(x_0) + \int_{x_0}^{z} F(t+L)w(t+L)\mathrm{d}t \\ &= v(x_0) + \int_{x_0}^{z} F(t)w(t+L)\mathrm{d}t, \\ w(z) &= v(x_0) + \int_{x_0}^{z} F(t)w(t)\mathrm{d}t. \end{aligned}$$

guarantees the periodicity of w also by the same criterion as (B.3).

Finally, since the all arguments above hold true for any ε with $0 < \varepsilon < d$, we obtain the conclusion. □